

Matične ćelije zubne pulpe i njihov potencijalni značaj u regenerativnoj medicini

DOI: 10.2298/SGS0803170T

Dental pulp stem cells - potential significance in regenerative medicine

Vera Todorović¹, Dejan Marković², Nadežda Milošević-Jovčić¹, Marijana Petakov¹, Bela Balint³, Miodrag Čolić⁴, Ana Milenković^{1,2}, Ivana Čolak¹, Vukoman Jokanović, Nebojša Nikolić²

¹Institut za medicinska istraživanja, Univerzitet u Beogradu

²Klinika za dečju i preventivnu stomatologiju, Stomatološki fakultet, Univerzitet u Beogradu

³Institut za transfuziju, Vojnomedicinska akademija u Beogradu

⁴Institut za medicinska istraživanja, Vojnomedicinska akademija u Beogradu

⁵Institut za nuklearna istraživanja Vinča, Univerzitet u Beogradu

¹ Institute for Medical Research, University of Belgrade

²Department of Preventive and Paediatric Dentistry, Faculty of Dentistry, University of Belgrade

³Institute for Tranfusion, Military Medical Academy in Belgrade

⁴Institute for Medical Research, Military Medical Academy in Belgrade

⁵Institute of Nuclear Sciences – Vinča, University of Belgrade

INFORMATIVNI RAD (IR) INFORMATIVE ARTICLE

KRATAK SADRŽAJ

Iz zubne pulpe su do danas izolovane tri populacije matičnih ćelija koje su označene kao matične ćelije zubne pulpe (engl. Dental Pulp Stem Cells, DPSC), matične ćelije iz eksfoliranih mlečnih zuba (engl. Stem Cells From Human Exfoliated Deciduous Teeth, SHED) i nezrele matične ćelije zubne pulpe (engl. Immature Dental Pulp Stem Cells, IDPC). Sve matične ćelije zubne pulpe su ektomezenhimalnog porekla i lokalizovane su u perivaskularnoj niši. One se lako i efikasno izoluju, visoko su proliferativne, klonogene, multipotentne, ispoljavaju visok stepen plasticiteta i i slične su mezenhimalnim matičnim ćelijama koštane srži (BMSC). U njima je pokazana visoka ekspresija gena alkalne fosfataze, proteina I matriksa dentina i dentin-sijalofosfoproteina. Takođe, istaknuta je važnost u ovim ćelijama ekspresije više gena koji kodiraju sintezu komponenti ekstracelularnog matriksa, molekula ćelijske adhezije, faktora rasta, transkripcionih faktora, gena prenosa ćelijskih signala, ćelijske komunikacije i metabolizma.

U uslovima in vitro ili in vivo ove ćelije mogu da se diferenciraju, s određenim međusobnim razlikama, u pravcu odontoblasta, hondročita, osteoblasta, adipocita, neurona/glijice, glatkih i skeletnih mišićnih ćelija, endotelnih ćelija i melanocita. U uslovima in vivo, nakon implantacije, pokazuju različit potencijal za formiranje dentina, ali i koštanog, masnog i nervnog tkiva.

SUMMARY

To date, three types of dental stem cells have been isolated: Dental Pulp Stem Cells (DPSC), Stem Cells From Human Exfoliated Deciduous Teeth (SHED) and Immature Dental Pulp Stem Cells (IDPC). These dental stem cells are considered as mesenchymal stem cells. They reside within the perivascular niche of dental pulp. They are highly proliferative, clonogenic, multipotent and are similar to mesenchymal Bone Marrow Stem Cells (BMSC). Also, they have high plasticity and can be easily isolated. The expressions of the alkaline phosphatase gene, dentin matrix protein 1 and dentin sialophosphoprotein are verified in these cells. Analyses of gene expression patterns indicated several genes which encode extracellular matrix components, cell adhesion molecules, growth factors and transcription regulators, cell signaling, cell communication or cell metabolism.

In both conditions, in vivo and in vitro, these cells have the ability to differentiate into odontoblasts, chondrocytes, osteoblasts, adipocytes, neurons, melanocytes, smooth and skeletal muscles and endothelial cells. In vivo, after implantation, they have shown potential to differentiate into dentin but also into tissues like bone, adipose or neural tissue. In general, DPSCs are considered to have antiinflammatory and immunomodulatory abilities.

Generalno se smatra da DPSC imaju anti-inflamatorno dejstvo i ispoljavaju imunom-odulatorni efekat. Takođe, dove do imunološke tolerancije ukoliko se implantiraju u alogena tkiva.

Sposobnost inhibicije proliferacije T limfocita ukazuje na njihovo imunosupresivno dejstvo. Matične ćelije zubne pulpe otvorile su nove perspektive u terapijskoj primeni ovih ćelija ne samo u regeneraciji dentina, tkiva periodoncijuma i koštano-zglobnog tkiva kraniofacijalne regije, već i u lečenju neurotraume, autoimunske oboljenja, infarkta miokarda, mišićne distrofije i oštećenja vezivnog tkiva.

Ključne reči: matične ćelije, zubna pulpa, regeneracija tkiva

Uvod

Matične ćelije su ključna podgrupa nespecijalizovanih ćelija organizma u vrlo ranom stadijumu razvića, koje u normalnim uslovima u datom tkivu mogu da se diferenciraju u različite tipove funkcionalno specijalizovanih zrelih ćelija. One igraju važnu ulogu u embrionalnom razviću i organogenesi (embrionalne i fetalne matične ćelije), kao i u tkivnoj homeostazi i regeneraciji (adultne matične ćelije)^{1,2}.

Matične ćelije se klasifikuju u odnosu na njihovu funkciju (normalne i kancerske matične ćelije) ili izvor izolacije (embrionalne, fetalne, matične ćelije iz krvi pupčanika i adultne matične ćelije). Postojanje humanih adultnih matičnih ćelija do danas je dokazano u koštanoj srži, perifernoj krvi, folikulu dlake, epitelu digestivnog trakta, skeletnom i srčanom mišiću, plućima, retini, mozgu, jetri, pankreasu, masnom tkivu, sinovijumu, periostijumu i zubu^{2,3,4,5}. Iako su u relativnom stanju mirovanja, ove ćelije su sposobne da snažno odgovore na tkivno oštećenje.

Najnovija istraživanja pokazuju da embrionalne i neke adultne matične ćelije mogu biti upotrebljene za regeneraciju dentina, tkiva periodoncijuma i rekonstrukciju kraniofacijalnih struktura.

Ne-hematopoetske mezenhimalne matične ćelije koštane srži (engl. Bone Marow Stromal Stem Cells, BMSC/MAPC), koje poseduju multipotentni potencijal i veliki plasticitet, mogu da se diferenciraju u osteoblaste, cementoblaste i ameloblaste, što otvara nove mogućnosti u regenerativnoj stomatologiji^{6,7}. Potrebno je istaći pionirski pokušaj i uspeh u kliničkoj upotrebi BMSC i plazme obogaćene trombocitima, upotrebo tehnologije tkivnog inženjeringu, u regeneraciji periodontalnog tkiva ljudi⁸. Takođe, dokazano je da kultivisane matične ćelije masnog tkiva pacova pomešane sa plazmom obogaćenom trombocitima i aplikovane lokalno životnjama sa periodontalnim defektom, indukuju duž alveolarne kosti stvaranje tkiva sličnog periodontalnom ligamentu⁹, pa se smatra da će u budućnosti zbog lake dostupnosti masnog tkiva dobijenog liposukcijom, matične ćelije masnog tkiva biti korisne u ćelijskoj terapiji oboljenja periodoncijuma.

After being grafted into allogenic tissues these cells are able to induce immunological tolerance.

Immunosuppressive effect is shown through the ability to inhibit proliferation of T lymphocytes.

Dental pulp stem cells open new perspectives in therapeutic use not only in dentin regeneration, periodontal tissues and skeletoarticular, tissues of craniofacial region but also in treatment of neurotrauma, autoimmune diseases, myocardial infarction, muscular dystrophy and connective tissue damages.

Keywords: stem cells, dental pulp, tissue regeneration

Introduction

Stem cells are unspecialized cells in early stage of development which have, in normal conditions, the ability to differentiate into a diverse range of specialized mature cells. They play important role in embryonic development and organogenesis (embryonic and fetal stem cells) as well as in homeostasis and tissue regeneration (adult stem cells)^{1,2}. Stem cells can be classified due to their function (normal and cancer stem cells) or potential source of isolation (embryonic, fetal, umbilical cord, adult stem cells). To date, adult stem cells have been isolated from many tissues including bone marrow, peripheral blood, hair follicle, digestive tract epithelium, skeletal and cord muscle, lungs, retina, brain, liver, pancreas, adipose tissue, synovium, periost and tooth^{2,3,4,5}. Although, these cells are in relatively inactive stage, they are capable of responding to tissue damage.

Recent advances in stem cell biology and gene therapy technology have provided the great potential of embryonic and some adult stem cells for use in dentin regeneration, periodontal tissue regeneration and reconstruction of cranifacial structures.

Non-hematopoietic mesenchymal BMSC posses multilineage potential and plasticity so they are able to differentiate into osteoblasts, cementoblasts and ameloblasts^{6,7}. These findings promise significant implications for regenerative dentistry. Also, recent study has demonstrated successful periodontal tissue regeneration with mesenchymal stem cells and platelet-rich plasma using tissue engineering technology⁸. Adipose Stem Cells (ASCs) were isolated from a rat, mixed with platelet-rich plasma obtained from inbred rats, and implanted into the periodontal tissue defect that had been generated in the test rats. Few weeks after implantation, a periodontal ligament-like structure was observed along with alveolar bone⁹. These observations suggest that ASCs can promote periodontal tissue regeneration in vivo. Because large amounts of human lipoaspirates are readily available, ASCs may be useful in future clinical cell-based therapy for periodontal diseases.

Smatra se da će embrionalne matične ćelije u budućnosti biti jedan od alternativnih izvora za rekonstrukciju kraniofacijalnih struktura, s obzirom da je dokazan njihov osteogeni potencijal u uslovima kultivisanja, dok se u uslovima in vivo ove ćelije diferenciraju u koštano tkivo u zubnoj alveoli pacova, bez pojave teratoma¹⁰.

Matične ćelije zubne pulpe lokalizovane su u perivaskularnoj niši i slične su BMSC. One su visoko proliferativne, klonogene, multipotentne i ispoljavaju visok stepen plasticiteta. Zbog lake dostupnosti, smatra se da će zubna pulpa, pored koštane srži, u budućnosti predstavljati najvažniji izvor adultnih multipotentnih mezenhimalnih matičnih ćelija, koje mogu da nađu široku primenu u regenerativnoj medicini^{11, 12}.

Izolacija, klasifikacija, molekulske i funkcionalne karakteristike matičnih ćelija zubne pulpe

Matične ćelije zuba mogu se izolovati iz: 1) zubnog epitela, 2) zubne pulpe, 3) periodontalnog ligamenta, 4) zubnog folikula i 5) apikalne papile zuba. Sve matične ćelije zuba, izuzev zubnih epitelnih matičnih ćelija koje se gube kod ljudi nakon izbijanja zuba i potiču od ektoderma, su ektomezenhimalnog porekla i potiču iz nervnog grebena. Osnovni potencijal diferencijacije matičnih ćelija zuba leži u njihovoj sposobnosti da formiraju dentin i tkiva periodoncijuma¹³.

Veoma je kompleksna karakterizacija različitih matičnih ćelija zuba na osnovu ekspresije površinskih proteinskih markera, kao i genske ekspresije (šema 1, tabela 1)^{13, 14}.


In recent study, the osteogenic potential of Embryonic Stem Cells (ESCs), using in vitro culture conditions was assessed. In vivo differentiation showed formation of osteogenic structure in the tooth sockets without evidence of teratomas¹⁰. These data suggest that pluripotent ESCs can serve as an alternative source for the reconstruction of craniofacial structures.

Dental pulp stem cells reside within the perivasculär niche of dental pulp and are considered as mesenchymal stem cells. They are highly proliferative, clonogenic, multipotent and also have high plasticity. Extraction of stem cells from pulp tissue is highly efficiency. The easy management of dental pulp stem cells make them, beside bone marrow stem cells, feasible for use in clinical trials on human patients and regenerative medicine^{11, 12}.

Isolation, classification, molecular and functional characteristics of dental pulp stem cells

Dental tissues which have been identified as accessible sources of stem cells are: 1) dental epithelium, 2) dental papilla, 3) periodontal ligament, and 4) dental follicle. All dental stem cells are neural crest-derived ectomesenchymal cells, with the exception of those from dental epithelium which are ectodermal cells. Constructing complex structures like dentin and periodontal tissue, dental stem cell biology might provide meaningful insights into the development of dental tissues and cellular differentiation processes¹³.

Characterization of diverse types of dental stem cells, due to expression of surface protein markers as well as gene expression, is very complexed (figure 1, table 1)^{13, 14}.



DPSC-Dental Pulp Stem Cell; DFPC-Dental Follicular Precursor Cell; SCAP-Stem Cell of Apical Papilla; PDLSC-Periodontal Ligament Stem Cell

Šema 1. Najvažniji površinski ćelijski markeri matičnih ćelija zuba*

* po Morsczeck et al., Clin Oral Invest 2008; 12:113-118

Antigen	DPSC*	SHED	PDLSC	BMSC
CD14	-	-	-	-
CD34	-	-	-	-
CD44	++	++	++	++
CD45	-	-	-	-
CD106	+	+/-	+/-	++
CD146	++/+-	++/+-	++/+-	++/+-
3G5	+/-	+/-	+/-	+/-
Stro-1	++/+-	++/+-	++/+-	++/+-
α-glatkomišićni aktin	++/-	++/-	++/-	++/+-
kolagen tip-I	++	++	++	++
kolagen tip-III	++/+	++/+-	++/+-	++/+
alkalna fosfataza	++/+-	++/+-	++/+	++/+-
osteokalcin	++/+	++/+-	++/-	+/-
osteonektin	++/+	++/+	++/+	++/+
osteopontin	+/-	+/-	+/-	+/-
sialoprotein kosti	-	-	-	-
skleraksis	+	+	++	+
sialofosfoprotein dentina	-	-	-	-

*DPSC-Dental Pulp Stem Cells; SHED-Stem Cells From Human Exfoliated Deciduous Teeth; PDLSC-Periodontal Ligament Stem Cells; (++) jaka ekspresija; (+) slaba ekspresija; (-) negativno; (/) subpopulacija

Tabela 1. Profil ekspresije proteina ili genske ekspresije u nekim matičnim ćelijama zuba u uslovima kultivisanja in vitro i odnos prema BMSC

Medutim, neki površinski proteinski markeri, kao što su Stro-1 i CD73 su ubikvitarno eksprimirani na svim matičnim i prekursorskim ćelijama zuba¹³. Posebna populacija matičnih ćelija zubne pulpe označena kao nezrele matične ćelije zubne pulpe (engl. Immature Dental Pulp Stem Cells, IDPSC) eksprimira markere embrionalnih matičnih ćelija Oct-4, Nanog, SSEA-3, SSEA-4, TRA-1-60 i TRA-1-81. Prepostavlja se da su ove ćelije u stvari pluripotentni prekursori druge dve populacije matičnih ćelija zubne pulpe označenih kao matične ćelije zubne pulpe u užem smislu reči (engl. Dental Pulp Stem Cells, DPSC) i matičnih ćelija zubne pulpe izolovanih iz eksfoliranih mlečnih zuba (engl. Stem Cells From Human Exfoliated Deciduous Teeth, SHED)¹⁵.

Regulacija homeostaze adultnih matičnih ćelija rezultat je suptilnog balansa između genetičkih i molekularnih ćelijskih mehanizama, spoljašnjih faktora iz lokalnih i sistemskih niša tela, kao i različitih puteva prenosa ćelijskih signala. Određivanjem profila genske ekspresije DPSC i funkcionalnom klasifikacijom tih gena ustanovljena je, pre svega, visoka ekspresija gena alkalne fosfataze, proteina 1 matriksa dentina (DMP1) i dentin-sijalofosfoproteina. Takođe, ukazano je na važnost u ovim ćelijama ekspresije više gena koji kodiraju sintezu komponenti ekstracelularnog matriksa, molekula ćelijske adhezije, faktora rasta i transkripcionih faktora. Funkcionalna analiza i analiza klasteriranja ukazala je važnost ekspresije gena prenosa ćelijskih signala, ćelijske komunikacije i ćelijskog metabolizma (tabela 2)¹⁴.

Some surface proteins, like Stro-1 and CD73, are ubiquitously expressed by all dental stem or precursor cells¹³. Population of Immature Dental Pulp Stem Cells (IDPSC) express embryonic stem cell markers Oct-4, Nanog, SSEA-3, SSEA-4, TRA-1-60 and TRA-1-81. It is assumed that these cells are pluripotent precursors to the other two stem cell populations known as Dental Pulp Stem Cells (DPSC) and Stem Cells From Human Exfoliated Deciduous Teeth (SHED)¹⁵.

To maintain homeostasis, a balance between genetic and molecular cell mechanism, extrinsic signals from the microenvironment (called niche) in which stem cells reside and signaling pathway in regulation of stem cell properties, is essential. Determining the profile of gene expression of dental pulp stem cells and functional classification of these genes, the expressions of the alkaline phosphatase gene, dentin matrix protein 1, and dentinsialophosphoprotein are verified and consequently the high expressions of these genes are discovered. Analyses of gene expression patterns indicated several genes which encode extracellular matrix components, cell adhesion molecules, growth factors, and transcription regulators. Functional and clustering analyses of differences in gene expression levels revealed cell signaling, cell communication, or cell metabolism (table 2)¹⁴.

Glavna kategorija gena	Podkategorija gena	hBMSC (I)*		hDPSC		hDPSC (I)		hOMC		hDPSC (I) (18-24)	
		↑	↓	↑	↓	↑	↓	↑	↓	↑	↓
deoba ćelije	DNK sinteza/replikacija	8	7	4	9	7	6	12	2	2	
	apoptoza	15	6	39	20	31	16	39	12	11	6
	ćelijski ciklus	25	8	27	31	39	15	68	7	11	14
	hromozom	6	5	4	9	5	3	9	1	1	3
	ukupno	54	26	74	69	82	40	128	22	25	23
signalni putevi ćelije ili ćelijska komunikacija	ćelijska athezija	16	14	33	22	23	24	32	26	10	4
	kanali/transport proteina	6	9	13	11	18	14	18	8	7	6
	faktori rasta/citokini	15	9	13	8	23	6	30	7	16	3
	metabolizam	18	12	17	20	28	14	42	8	10	4
	modifikacija proteina	5	8	20	18	23	3	12	10	14	9
struktura ćelije / motilitet	receptori	59	32	78	53	65	44	81	62	28	16
	Ukupno	119	84	174	132	180	105	215	121	85	42
	citoskelet	4	3	5	7	6	5	11	2	2	5
	ekstracelularni matriks	30	11	17	25	15	31	15	31	5	7
	protein vezan za mikrotubule	1		7		5		14		3	
ćelija / odbrana organizma	ukupno	35	14	29	32	26	36	40	33	7	15
	/ homeostaza (odgovor na stres)	41	14	40	45	61	25	56	39	28	16
	imunologija (imuni odgovor)	46	10	41	30	53	17	51	46	24	11
gen ekspresija proteina	ukupno	87	24	81	75	114	42	107	85	52	27
	/ RNK sinteza (transkripcioni faktor)	31	16	62	49	52	21	55	31	35	13
	sinteza proteina (vezivanje proteina)	4	3	3	7	6	2	5	2	2	5
metabolizam	ukupno	35	19	65	56	58	23	60	33	37	18
	aminokiselina	5	1	7	4	9	5	3	5	7	1
	energija/TCA ciklus	1	1	3		3		4			
	lipid	21	4	27	7	33	6	21	6	13	1
	nukleotid	2	2	4	7	6	4	21	8	4	8
	modifikacija proteina	13	12	41	32	20	3	13	21	19	11
	transport	45	19	50	39	83	32	62	35	39	12
	ukupno	87	39	132	89	154	50	124	75	82	33
	UKUPNO	417	206	555	453	614	296	674	369	288	158

*hBMSC (I)-humane mezenhimalne matične ćelije koštane srži u osteoidnuktivnoj sredini 18. dana kultivacije; hDPSC (I)-humane matične ćelije zubne pulpe u osteoinsuktivnoj sredini 18. dana kultivacije; hOMC-humane ćelije oralne mukoze; hDPSC (I) (18-24)- humane matične ćelije zubne pulpe u osteoinsuktivnoj sredini 18-24 dana kultivacije

Tabela 2. Funkcionalna klasifikacija gena hDPSC

Što se tiče razlika u genskoj ekspresiji između MAPC/BMSC i DPSC, one se ogledaju pre svega u povećanoj ekspresiji kod DPSC gena koji kontrolišu ćelijski ciklus, naročito gena ciklin-zavisne kinaze-6 koja je aktivator ćelijskog ciklusa, što objašnjava višu stopu proliferacije DPSC u odnosu na MAPC/BMSC^{14, 16}. Put prenosa signala Wnt/beta-katenin¹⁷, koji je ključan za samoobnavljanje matičnih ćelija, kao i put Notch¹⁸, negativno regulišu diferencijaciju in vitro DPSC u odontoblaste, dok induktivni efekat imaju FGF2 i TGFbeta¹⁹. U poslednje vreme poznato je da je hipoksija jedan od ključnih faktora u održavanju nediferenciranog stanja i plasticiteta adultnih matičnih ćelija²⁰. Međutim, uslovi kultivisanja koji imitiraju ishemiju (hipoksiju i hipoglikemiju) nepovoljno utiču na preživljavanje i diferencijaciju DPSC²¹. Bolje poznavanje ekspresije površinskih markera i gena matičnih ćelija zubne pulpe, puteva prenosa signala, i njihov odnos prema ćelijskoj diferencijaciji i plasticitetu, svakako će u budućnosti doprineti razvoju terapije bazirane na primeni ovih ćelija u humanoj medicini i stomatologiji.

Matične ćelije zubne pulpe imaju sve osobine potrebne za uspešnu terapijsku primenu: 1) lako su dostupne za izolaciju; 2) izolacija je veoma efikasna; 3) imaju multipotentni potencijal; 4) pokazuju interakciju sa biomaterijalima koji upotrebljeni kao matrice indukuju intenzivnu proliferaciju ovih ćelija; 5) dugovečne su i 6) podležu uspešnoj krioprotekciji, slično drugim matičnim ćelijama^{11, 22}. Podatak da vijabilne matične ćelije zubne pulpe mogu da se izoluju i nakon 5 dana od ekstrakcije zuba, kao i da ih je moguće izolovati i kultivisati iz intaktnih zuba koji su prethodno krioprezervirani, ukazuje da su potrebni minimalni zahtevi za stvaranje banke ovih ćelija u cilju buduće terapijske primene¹².

Studijama in vitro je dokazano da matične ćelije zubne pulpe imaju izrazit plasticitet, tj. sposobnost da se diferenciraju u ćelije različite od ćelija prisutnih na mestu odakle potiču, a najnoviji radovi pružaju dokaze o njihovom plasticitetu i u uslovima in vivo. U uslovima in vitro ili in vivo ove ćelije mogu da se diferenciraju, s određenim međusobnim razlikama, u pravcu odontoblasta, hondročita, osteoblasta, adipocita, neurona/glije, glatkih i skeletnih mišićnih ćelija, endotelnih ćelija i melanocita^{15, 16, 23-29}. U uslovima in vivo, nakon implantacije, pokazuju različit potencijal za formiranje dentina, ali i koštanog i masnog tkiva³⁰.

Generalno se smatra da DPSC, slično drugim mezenhimalnim matičnim ćelijama, imaju anti-inflamatorno dejstvo. One ispoljavaju imunomodulatorni efekat jer mogu biti uključene u imunski odgovor u toku infekcije zubne pulpe, a verovatno i periodoncijuma, koji je veoma kompleksan^{31, 32}, putem aktivacije NF- κ B³³. Takođe, dovode do imunološke tolerancije ukoliko se implantiraju u alogena tkiva¹¹.

Comparing the gene expression profiles of human dental pulp stem cells to bone marrow stromal stem cells a few differentially expressed genes, including cell cycle activator cyclin-dependent kinase 6, were highly expressed in DPSCs. This explains the higher proliferation rate of dental pulp stem cells than bone marrow stromal stem cells^{14, 16}. Wnt/beta-catenin, important for stem cells self-renewal¹⁷, and Notch signaling¹⁸ can inhibit the odontoblastic differentiation, while two crucial growth factors, FGF2 and TGFbeta¹⁹, have inductive effects on the odontoblastic differentiation of human dental pulp stem cells in vitro. Although differentiated and undifferentiated cells can be exposed to ischemic conditions in cases of injury or inflammation, the effects of ischemia on cell survival and differentiation have not been well characterized²⁰. These data showed that the ischemic conditions have similar detrimental influence on both undifferentiated and differentiated porcine Dental Pulp-derived Cells (pDPCs), and affect differentiation status of pDPCs²¹. Dental pulp stem cells have all characteristics for successful therapeutic use: 1) access to the collection site of these cells is easy, 2) extraction of stem cells from pulp tissue is highly efficiency; 3) they have an extensive differentiation ability, 4) demonstrated interactivity with biomaterials makes them ideal for tissue reconstruction, 5) they have a long lifespan, 6) they can be safely cryopreserved^{11, 22}. Recent studies indicate that DPSC isolation is feasible for at least 5 days after tooth extraction. Further, the recovery of viable DPSC after cryopreservation of intact teeth suggests that minimal processing may be needed for the banking of samples with no immediate plans for expansion and use¹².

An increasing number of investigations supports that stem cells have the potential to differentiate into matured cell types beyond their origin, a property defined as plasticity. Previously, the plasticity of DPSCs has been confirmed by culturing cells in lineage-specific media in vitro. In recent studies, it has been confirmed in vivo conditions as well. In both conditions, in vivo and in vitro, these cells have the ability to differentiate into odontoblasts, chondrocytes, osteoblasts, adipocytes, neurons, melanocytes, smooth and skeletal muscles and endothelial cells^{15, 16, 23-29}. In vivo, after implantation, they have shown potential to differentiate into dentin but also into bone or adipose tissue³⁰.

In general, DPSCs are considered to have anti-inflammatory and immunomodulatory abilities because they may be involved in immune responses during pulpal infection, and probably periodontium as well^{31, 32}, through activating NF- κ B³³. After being grafted into allogenic tissues these cells are able to induce immunological tolerance¹¹.

Sposobnost inhibicije proliferacije T limfocita ukazuje na njihovo imunosupresivno dejstvo i moguću primenu kod autoimunskih bolesti³⁴.

Različite vrste matičnih ćelija zubne pulpe i njihov potencijalni značaj u regenerativnoj medicini i stomatologiji

Sredinom devedesetih godina XX. veka došlo je do uspešne izolacije prekursorskih ćelija zubne pulpe³⁵. Kasnije su matične ćelije izolovane iz zubne pulpe kutnjaka i nazvane DPSC²³, kao i iz eksfoliranih mlečnih zuba-SHED³⁶ i IDPC¹⁵.

DPSC

Ove ćelije su do sada izdvojene iz umnjaka, mlečnih sekutića i prekobrojnih zuba^{16, 23, 37}. Kao što je već iustaknuto, ispoljavaju slične karakteristike kao BMSC³⁸.

Prva istraživanja na DPSC izolovanim iz umnjaka su pokazala da se ove ćelije diferenciraju u odontoblaste, adipocite i u neuronima-slične ćelije i da ta diferencijacija podseća na embrionalnu ontologiju kranijalnih ćelija nervnog grebena. Istovremeno je pokazano i da imaju neke karakteristike identične sa osteoblastima²⁴. Međutim, najnovija istraživanja pokazuju da humane (h) DPSC imaju daleko veći potencijal diferencijacije, s obzirom da mogu da se diferenciraju u ćelije mezenhimalnog porekla (odontoblaste, osteoblaste, adipocite, hondrocite i poprečnoprugaste mišićne ćelije) ali i u melanocite (ćelije koje nisu porekla mezenhima, već potiču od ćelija nervnog grebena)²⁶. Stoga se pretpostavlja da se u kulturama hDPSC nalaze i ćelije koje bi mogle biti multipotentne matične ćelije nervnog grebena.

Pokazano je da humane DPSC izolovane iz mlečnih zuba, koje u uslovima in vitro pokazuju osteogeni, adipogeni i miogeni potencijal diferencijacije, transplantirane pacovima sa velikim defektom parijetalne regije, dovode do stvaranja nove kosti nakon 1 meseca od načinjene lezije, bez upotrebe imunosupresivne terapije²⁵. To daje nadu da će se ove ćelije u budućnosti moći upotrebiti u rekonstruktivnoj hirurgiji kraniofacijalne regije.

Matične ćelije označene kao SBP-DPSC su multipotentna subpopulacija DPSC, koja je sposobna da se diferencira u osteoblaste i da sintetiši trodimenzionalnu lamelarnu kost in vitro. Ukoliko se SBP-DPSC ili koštani model dobijen iz njih u uslovima in vitro transplantira pacovu sa deficitom imunološkog sistema stvorice se tkivna struktura sa integralnim krvnim sudovima, slična pravoj kosti odraslih ljudi^{28,29}. Dokazano je da čak i nakon 2 godine od krioprezervacije SBP-DPSC zadržavaju sposobnost diferencijacije, proliferacije i sinteze koštanog tkiva²⁹.

Immunosuppressive effect and possible therapeutic use in autoimmune diseases is shown through the ability to inhibit proliferation of T lymphocytes³⁴.

Different types of dental pulp stem cells - potential significance in regenerative medicine and dentistry

In the last decade of the twentieth century, precursor cells of dental pulp were successfully isolated³⁵. Furthermore, DPSCs²³ were isolated from dental pulp of human adult third molar as well as from human exfoliated deciduous teeth³⁶ and immature dental pulp¹⁵.

DPSC

To date, these cells have been isolated from third molars, decidual incisors and supernumerary teeth^{16, 23, 37}. DPSC have similar characteristics to Bone Marrow Stem Cells (BMSC)³⁸.

DPSCs derived from third molars were also found to be capable of differentiating into odontoblasts, adipocytes and neural-like cells and that differentiation reminiscent of cranial neural crest (CNC) cells embryonic ontology. At the same time, it is found that DPSCs have some characteristics identical to osteoblasts²⁴. It is demonstrated that human dental pulp contains self renewing human Dental Pulp Stem Cells (hDPSCs) capable of differentiating into mesenchymal-derived odontoblasts, osteoblasts, adipocytes, chondrocytes and striated muscle, and interestingly, also into non-mesenchymal, neural crest- derived melanocytes²⁶. Furthermore, this study showed that hDPSC cultures include cells with traits attributed to multipotent stem cells, and provide evidence that these might be multipotent neural crest stem cells.

Human dental pulp stem cells, were isolated from deciduous teeth, to reconstruct large-sized cranial bone defects in nonimmunosuppressed rats. They showed osteogenic, adipogenic, and myogenic in vitro differentiation²⁵. These findings suggest that hDPSC is an additional cell resource for correcting large cranial defects in rats and constitutes a promising model for reconstruction of human large cranial defects in craniofacial surgery.

Stromal stem cells from human dental pulp (SBP-DPSCs) are multipotent stem cells able to differentiate into osteoblasts, which synthesize three-dimensional woven bone tissue chips in vitro. When either SBP-DPSCs or bone chips obtained in vitro were transplanted into immunocompromised rats, they generated a tissue structure with an integral blood supply similar to that of human adult bone^{28,29}. After storage for 2 years, it is found that stem cells are still capable of differentiation, and that their differentiated cytotypes proliferate and produce woven bone tissue²⁹.

Subpopulaciju DPSC, označenu kao SP-DPSC imaju veliki angiogenetski potencijal. Ukoliko se miševima sa ishemijom prednje šape lokalno transplantiraju ove ćelije doći će do njihovog uspešnog kalemjenja, povećanja gustine kapilarne mreže i bolje prokrvljenosti ishemijске zone, a kondicionirani medijum od ovih ćelija pokazuje mitogenu i anti-apoptotsku aktivnost na endotelne ćelije humane umbilikalne vene³⁹.

Stvaranje in vivo tkiva sličnog zubnoj pulpi, nakon subkutane aplikacije mišu trijade DPSC, matice od kolagena i DMP1, ukazuje na veliki potencijalni značaj ovih ćelija u tkivnom inženjeringu⁴⁰.

SHED

Matične ćelije iz eksfoliranih mlečnih zuba su sposobne za diferencijaciju u odontoblaste, adipocite i nervne ćelije. U uslovima in vivo one indukuju formiranje kosti i produkciju dentina; nakon transplantacije u miša sa kompromitovanim imunološkim sistemom migriraju u možak i sposobne su za preživljavanje 36. Studije in vivo su takođe pokazale da se humane SHED diferenciraju u odontoblaste i endotelne ćelije i da stoga eksfolirani mlečni zubi predstavljaju vijabilni izvor matičnih ćelija za tkivni inženjerинг Zubne pulpe. Naime, ukoliko se hSHED oblože biodegradibilnim materijalom pripmljenim sa presecima normalnog zuba i kao takve implantiraju mišu sa imunološkim deficitom, oformiće se tkivo koje u histološkom pogledu i u pogledu ćelijskog sastava veoma podseća na zubnu pulpu⁴¹.

IDPC

Radi se o najprimitivnijoj populaciji matičnih ćelija Zubne pulpe, s obzirom da je pokazano da eksprimiraju markere embrionalnih matičnih ćelija. U toku kultivisanja kroz 25 pasaža, u toku 4 meseca, one zadržavaju normalni kariotip i stopu ekspanzije karakterističnu za matične ćelije. U hemijski definisanim uslovima kultivisanja in vitro, IDPC se diferenciraju u glatke i poprečnoprugaste mišićne ćelije, neurone, hondrocite i osteoblaste. U uslovima in vivo, ukoliko se transplantiraju miševima sa imunološkim deficitom, dobro se kaleme u različitim tkivima 15. Aplikacija hIDPC bez primene imunosupresivne terapije, pokazala se uspešnom u kontroli mišićne distrofije kod zlatnih retrivera (najboljeg animalnog model za Duchenne-ovu mišićnu distrofiju kod ljudi), pri čemu je sistemska aplikacija ovih matičnih ćelija bila bolji izbor od lokalne aplikacije⁴².

Zaključak

Matične ćelije Zubne pulpe otvorile su nove perspektive u terapijskoj primeni ovih ćelija ne samo u regeneraciji dentina, tkiva periodoncijuma i koštano-zglobnog tkiva kraniofacijalne regije, već i u lečenju neurotraume, infarkta miokarda, mišićne distrofije i oštećenja vezivnog tkiva.

Side population (SP) dental pulp stem cells, SP-DPSC, has high angiogenetic potential. In models of mouse hindlimb ischemia, local transplantation of one subfraction of SP cells resulted in successful engraftment and an increase in the blood flow including high density of capillary formation. Conditioned medium from this subfraction showed the mitogenic and anti-apoptotic activity on human umbilical vein endothelial cells³⁹.

In vivo generation of dental pulp-like tissue by using the triad of dental pulp stem cells, a collagen scaffold, and dentin matrix protein 1 after subcutaneous transplantation in mice, might lead to hard tissue formation. This finding indicate that these cells might have high potential significance in tissue engineering⁴⁰.

SHED

SHED were identified to be a population capable of differentiating into a variety of cell types including neural cells, adipocytes, and odontoblasts. After in vivo transplantation, SHED were found to be able to induce bone formation, generate dentin and survive in mouse brain along with expression of neural markers 36. SHED differentiated into odontoblast-like cells in vivo and in endothelial-like cells. SHED seeded in biodegradable scaffolds prepared within human tooth slices and transplanted into immunodeficient mice result in tissue with architecture and cellularity that closely resemble those of a physiologic dental pulp. This work suggests that exfoliated deciduous teeth constitute a viable source of stem cells for dental pulp tissue engineering⁴¹.

IDPC

The population of immature dental pulp stem cells is dental pulp stem cell population which expresses embryonic stem cell markers during at least 25 passages while maintaining the normal karyotype and the rate of expansion characteristic of stem cells. Moreover, in vitro these cells can be induced to undergo uniform differentiation into smooth and skeletal muscles, neurons, cartilage, and bone under chemically defined culture conditions. After in vivo transplantation of these cells into immunocompromised mice, they showed dense engraftment in various tissues 15. Human Immature Dental Pulp Stem Cells (hIDPSC) were transplanted, without any immunosuppression, into golden retriever muscular dystrophy dogs, who represent the best available animal model for therapeutic trials aiming at the future treatment of human Duchenne muscular dystrophy. Data from this trial suggested that systemic multiple deliveries seemed more effective than local injections. These findings open important avenues for further researches⁴².

Conclusion

Dental pulp stem cells open new perspectives in therapeutic use not only in dentin regeneration, periodontal tissues and skeletoarticular tissues of craniofacial region but also in treatment of neurotrauma, autoimmune diseases, myocardial infarction, muscular dystrophy and connective tissue damages.

Literatura / References

1. Todorović V, Nikolić I, Glibetić M, Balint B. Humane embrionalne matične ćelije – dosadašnja saznanja. *Anestezija, reanimacija, transfuzija* 2006; 34:109-28
2. Todorović V, Nikolić RI. Stem ćelije, kloniranje sisara. U: Ivan R. Nikolić, urednik i ilustrator. Embriologija čoveka, tekst i atlas (treća izdanje). Beograd: Data Status, 2007: 45-70
3. Malanchi I, Peinado H, Kassen D, et al. Cutaneus cancer stem cell maintenance is dependent on catenin signalling. *Nature* 2008; 452:650-3
4. Brašanac D, Boričić I, Todorović V, Tomanović N, Radojević S. Cyclin A and β -catenin expression in actinic keratosis, Bowen's disease and invasive squamous cell carcinoma of the skin. *Br J Dermatology* 2005; 153:1166-75
5. Vats A, Bielby RC, Tolley NS, Nerem R, Polak JM. Stem cells. *Lancet* 2005; 366:592-602
6. Maria OM, Khosravi R, Mezey E, Tran SD. Cells from bone marrow that evolve into oral tissues and their clinical applications. *Oral Dis* 2007; 13:11-16
7. Ueda M, Yamada Y, Ozawa R, Okazaki Y. Clinical case reports of injectable tissue-engineered bone for alveolar augmentation with simultaneous implant placement. *Int J Periodontics Restorative Dent* 2005; 25:129-37
8. Yamada Y, Ueda M, Hibi H, Baba S. A novel approach to periodontal tissue regeneration with mesenchymal stem cells and platelet-rich plasma using tissue engineering technology: A clinical case report. *Int J Periodontics Restorative Dent* 2006; 26:363-9
9. Tobita M, Uysal AC, Ogawa R, Hyakusoku H, Mizuno H. Periodontal tissue regeneration with adipose-derived stem cells. *Tissue Eng Part A* 2008; 14:945-53
10. Kang HK, Roh S, Lee G, Hong SD, Kang H, Min BM. Osteogenic potential of embryonic stem cells in tooth sockets. *Int J Mol Med*. 2008;21:539-44
11. d'Aquino R, Papaccio G, Laino G, Graziano A. Dental pulp stem cells: a promising tool for bone regeneration. *Stem Cell Rev* 2008;4:21-6
12. Perry BC, Zhou D, Wu X, et al. Collection, cryopreservation, and characterization of human dental pulp-derived mesenchymal stem cells for banking and clinical use. *Tissue Eng Part C Methods* 2008;14:149-56
13. Morsczeck C, Schmalz G, Reichert TE, Volner F, Galler K, Driemel O. Somatic stem cells for regenerative dentistry. *Clin Oral Invest* 2008; 12:113-8
14. Yamada Y, Fujimoto A, Ito A, Yoshimi R, Ueda M. Cluster analysis and gene expression profiles: a cDNA microarray system-based comparison between human dental pulp stem cells (hDPSCs) and human mesenchymal stem cells (hMSCs) for tissue engineering cell therapy. *Biomaterials* 2006; 27:3766-81
15. Kerkis I, Kerkis A, Dozortsev D, et al. Isolation and characterization of a population of immature dental pulp stem cells expressing OCT-4 and other embryonic stem cell markers. *Cells Tissues Organs* 2006; 184:105-16
16. Shi S, Gronthos S. Perivascular niche of postnatal mesenchimal stem cells in human bone marrow and dental pulp. *J Bone Miner Res* 2003; 18:696-704
17. Scheller EL, Chang J, Wang CY. Wnt/beta-catenin inhibits dental pulp stem cell differentiation. *J Dent Res* 2008;87:126-30
18. Zhang C, Chang J, Sonoyama W, Shi S, Wang CY. Inhibition of human dental pulp stem cell differentiation by Notch signaling. *J Dent Res* 2008;87:250-5
19. He H, Yu J, Liu Y, et al. Effects of FGF2 and TGFbeta1 on the differentiation of human dental pulp stem cells in vitro. *Cell Biol Int* 2008;32:827-34
20. Kovacevic-Filipovic M, Petakov M, Hermitte F, et al. Interleukin-6 (IL-6) and low O-2 concentration (1%) synergize to improve the mainteance of hematopoietic stem cells (Pre-CSF). *J Cell Physiol* 2007; 212:68-75
21. Agata H, Kagami H, Watanabe N, Ueda M. Effect of ischemic culture contitions on the survival and differentiation of porcine dental pulp-derived cells. *Differentiation* 2008; DOI: 10.1111/j.1432-0436.2008.00282.x
22. Škorić D, Balint B, Petakov M, Sindjić M, Rodić P. Collection strategies and cryopreservation of umbilical cord blood. *Transfusion Med* 2007; 17:107-13
23. Gronthos S, Mancani M, Brahim J, Robey PG, Shi S. Postnatal human dental pulp stem cells (DPSCs) in vitro and in vivo. *Proc Natl Acad Sci* 2000; 97:13625-30
24. Gronthos S, Brahim J, Li W, et al. Stem cell properties of human pulp stem cells. *J Dent Res* 2002; 81:531-5
25. de Mendonça Costa A, Bueno DF, Martins MT, et al. Reconstruction of large cranial defects in nonimmunosuppressed experimental design with human dental pulp stem cells. *J Craniofac Surg* 2008; 19:204-10

26. Stevens A, Zuliani T, Olejnik C, et al. Human dental pulp stem cells differentiate into neural crest-derived melanocytes and have label-retaining and sphere-forming abilities. *Stem Cells Dev* 2008; Mart 25. [Epub ahead of print]
27. Arthur A, Rychkov G, Shi S, Koblar SA, Gronthos S. Adult human dental pulp stem cells differentiate toward functionally active neurons under appropriate environmental cues. *Stem Cells* 2008; 26:1787-95
28. d'Aquino R, Graziano A, Sampaolesi M, et al. Human postnatal dental pulp cells co-differentiate into osteoblasts and endotheliocytes: a pivotal synergy leading to adult bone tissue formation. *Cell Death Differ* 2007; 14:1162-71
29. Papaccio G, Graziano A, d'Aquino R, et al. Long-term cryopreservation of dental pulp stem cells (SBP-DPSCs) and their differentiated osteoblasts: a cell source for tissue repair. *J Cell Physiol* 2006; 208:319-25
30. Zhang W, Walboomers XF, Van Kuppevelt TH, et al. In vivo evaluation of human dental pulp stem cells differentiated towards multiple lineages. *J Tissue Eng Regen Med* 2008; 2:117-25
31. Colic M, Lukic A, Vucevic D, et al. Correlation between phenotypic characteristic of mononuclear cells isolated from human periapical lesions and their in vitro production of Th1 and Th2 cytokines. *Arch Oral Biol* 2006; 51:1120-30
32. Stefanović G, Marković D, Ilić V, Brajović G, Petrović S, Milošević-Jovčić N. Hypogalactosylation of salivary and gingival fluid IgG in patients with advanced periodontitis. *J Periodontol* 2006; 77:1887-93
33. Chang J, Zhang C, Tani-Ishii N, Shi S, Wang C-Y. NF- \square B activation in human dental pulp stem cells by TNF and LPS. *J Dent Res* 2005; 84:994-8
34. Pierdomenico L, Bonsi L, Calvitti M, et al. Multipotent mesenchymal stem cells with immunosuppressive activity can be easily isolated from dental pulp. *Transplantation* 2005; 80:836-42
35. Stanislawski L, Carreau JP, Pouchelet M, Chen ZN, Goldberg M. In vitro culture of human dental pulp cells: some aspects of cells emerging early from the explant. *Clin Oral Investig* 1997; 1:131-40
36. Miura M, Gronthos S, Zhao M, et al. SHED: Stem cells from human exfoliated deciduous teeth. *Proc Natl Acad Sci USA* 2003; 100:5807-12
37. Huang AH, Chen YK, Lin LM, Shieh TY, Chan AW. Isolation and characterization of dental pulp stem cells from a supernumerary tooth. *J Oral Pathol Med* 2008; Mart 5. [Epub ahead of print]
38. Shi S, Rhobey PG, Gronthos S. Comparison of human dental pulp and bone marrow stromal cells by cDNA microarray analysis. *Bone* 2001; 29:532-9
39. Johara K, Zheng L, Wake H, et al. A novel stem cell source for vasculogenesis in ischemia: subfraction of side population cells from dental pulp. *Stem Cells* 2008; Jun 26. [Epub ahead of print]
40. Prescott RS, Alsanea R, Fayad MI, et al. In vivo generation of dental pulp-like tissue by using dental pulp stem cells, a collagen scaffold, and dentin matrix protein 1 after subcutaneous transplantation in mice. *J Endod* 2008; 34:421-6
41. Cordeiro MM, Dong Z, Kaneko T, et al. Dental pulp tissue engineering with stem cells from exfoliated deciduous teeth. *J Endod* 2008; 34:962-9
42. Kerkis I, Ambrosio CE, Kerkis A, et al Early transplantation of human immature dental pulp stem cells from baby teeth to golden retriever muscular dystrophy (GRMD) dogs: local or systemic? *J Transl Med* 2008; 6:35

Adresa za korespondenciju

dr sci. med. Vera Todorović, naučni savetnik
 Institut za medicinska istraživanja,
 Univerzitet u Beogradu
 Dr Subotića 4, 11129 Beograd, PF 129
 e-mail: verat@imi.bg.ac.yu
 Tel:011/2685-788/L-111 ili 113

Address for correspondence

Vera Todorovic, research professor
 Institute for medical research
 University of Belgrade
 Dr Subotica 4 11000 Belgrade, PO BOX 129
 e-mail: verat@imi.bg.ac.yu
 Tel:+381 11/2685-788/ L-111 ili 113