

ORIGINAL ARTICLE / ОРИГИНАЛНИ РАД

Prevalence of developmental dental anomalies in Serbian orthodontic patients

Evgenija Marković¹, Ana Vuković², Tamara Perić², Jovana Kuzmanović-Pficer³, Bojan Petrović⁴¹University of Belgrade, School of Dental Medicine, Clinic of Orthodontics, Belgrade, Serbia;²University of Belgrade, School of Dental Medicine, Clinic for Pediatric and Preventive Dentistry, Belgrade, Serbia;³University of Belgrade, School of Dental Medicine, Department of Medical Statistics and Informatics, Belgrade, Serbia;⁴University of Novi Sad, Faculty of Medicine, Department of Pediatric and Preventive Dentistry, Novi Sad, Serbia**SUMMARY****Introduction/Objective** The aim of this study was to evaluate the prevalence of developmental dental anomalies (DDA) in Serbian orthodontic patients.**Methods** The sample was composed of 1,001 panoramic radiographs of orthodontic patients, older than seven years, taken as a part of the initial diagnostic procedure at the Clinic of Orthodontics, School of Dental medicine in Belgrade. The DDA that could be diagnosed accurately on panoramic X-rays were documented. Descriptive analysis was used to determine prevalence and sex distribution of DDA. The Pearson χ^2 test and Fisher's exact test were used to compare number of affected teeth in males and females (level of significance was 95%).**Results** The prevalence of DDA in Serbian orthodontic patients was 34.8% (15.5% males and 19.3% females). Impactions were present in 16.5%, hypodontia in 12.9%, hyperdontia in 4.4%, microdontia in 2.9%, macrodontia in 1.8% and transposition in 0.8% of patients. Maxillary canines were the most frequently impacted teeth. Maxillary second molars were more prone to impaction in females ($p < 0.05$). Impacted incisors were more prevalent in maxilla, premolars, and second molars in mandible. The most commonly missing teeth were upper left second premolars. Mesiodens was the most frequently found supernumerary tooth.**Conclusion** We reported a high a rate of DDA in Serbian orthodontic patients, more in females than males. The most frequently observed DDA were impaction, tooth agenesis, hyperdontia, microdontia, macrodontia, and transposition. All investigated DDA were more frequently present in females, except hyperdontia. Current findings could offer a foundation for epidemiological studies on DDA prevalence.**Keywords:** developmental dental anomalies; orthodontics; hypodontia**INTRODUCTION**

Developmental dental anomalies (DDA) occur during the period of teeth development. The etiology is complex and multifactorial. It involves genetic and environmental influences, as well as variation in sex distribution. DDA are presented as irregularities in tooth number, size, shape, and structure, and altered teeth eruption. The complexity of tooth development is influenced by over 300 genes, mutations, and/or localized or generalized insults (trauma, infection, therapeutic irradiation, low birth weight, vitamin D deficiency, metabolic and hormonal disturbances, as well as nutrition and available space in the dental arch). The outcome of these influences could be the presence of isolated or combined DDA in a person [1, 2]. Persons with DDA tend to have orthodontic, functional, and esthetic problems. The early discovery and information of prevalence and association of dental abnormalities with sex and type of teeth are important information for dental practitioners.

Epidemiological studies investigating the prevalence of DDA have been conducted all

over the world with variation in results [3, 4, 5]. Only a few recent studies, mostly on a particular type of DDA, were done in Serbia. Authors investigated the prevalence of hypodontia in Serbian schoolchildren [6, 7]. Two studies reported on the prevalence of structural dental anomalies (amelogenesis imperfecta and molar-incisor hypomineralization [8, 9]). To the best of our knowledge, any other studies investigating more types of DDA in Serbian population have not been conducted.

The aim of this study was to evaluate the prevalence and sex distribution of developmental dental anomalies in Serbian orthodontic patients.

METHODS**Sample**

This retrospective cross-sectional study was comprised of 1,324 panoramic radiographs of patients older than seven years of age referred to the Clinic of Orthodontics, School of Dental

Received • Примљено:

July 1, 2019

Revised • Ревизија:

August 8, 2019

Accepted • Прихваћено:

August 22, 2019

Online first: September 13, 2019**Correspondence to:**Ana VUKOVIĆ
Clinic for Pediatric
and Preventive Dentistry
Dr Subotića 11
11000 Belgrade, Srbija
ana.vukovic@stomf.bg.ac.rs

Medicine, University of Belgrade from all over Serbia. Digital panoramic radiographs were taken as a part of the initial diagnostic examination in 2016. Only high-quality films of patients with no craniofacial abnormalities and syndromes associated with DDA (including cleft lip and palate), previously extracted permanent teeth, a trauma in the orofacial region, and previous orthodontic treatment with fixed appliances, were included in the sample. Consequently, a sample comprised of 1,001 panoramic radiographs (459 male and 542 female patients). Experienced orthodontist and pedodontist assessed the radiographs. Only tooth abnormalities that could be diagnosed precisely and solely on panoramic X-rays were documented. The DDA with a high probability of poor diagnosis without previous clinical examination and/or additional radiographs were excluded from the evaluation, such as:

- 1) Anomalies of tooth structure – hypomineralization, amelogenesis imperfecta, and molar-incisor hypomineralization (MIH);
- 2) Root deformation and number, concrescence and dilaceration;
- 3) Rotation.

Third molars were excluded from the evaluation due to the high incidence of variation in morphology, size, and position.

We evaluated panoramic radiographs for the following DDA:

- 1) Hypodontia – developmentally missing teeth (tooth agenesis) was diagnosed by counting present teeth when no sign of tooth formation existed. Oligodontia was defined when more than six teeth were missing;
- 2) Hyperdontia (supernumerary teeth) – additional teeth were present on the radiograph. They may be observed as teeth with normal size and shape, or with smaller size and atypical form;
- 3) Mesiodens – supernumerary tooth localized in the anterior region of maxilla;
- 4) Tooth transposition – two adjacent teeth changed their position partially or completely in dental arch [10];
- 5) Microdontia – teeth are smaller than average. Microdontia of maxillary lateral incisor was recorded when the maximum mesiodistal crown diameter was smaller compared to the same dimension of opposing mandibular lateral incisor in the same patient [11];
- 6) Macrodonia was referred to the tooth that was found to be immensely larger than the average one [12];
- 7) Impaction was defined in cases when physical barrier existed, and/or tooth had an orientation that prevented its emergence [13]. Canines were not evaluated for impaction in children younger than ten years of age due to the possibility of misdiagnosis.

Statistical analysis

The statistical analyses were performed using Statistical Package for Social Science, version 22.0 (IBM Corp., Armonk, NY, USA). The descriptive statistical analysis was used to evaluate the prevalence of DDA and sex distribution. Pearson's χ^2 test and Fisher's exact test were used to

compare number of teeth affected by anomalies in males and females. The level of significance was set at $p < 0.05$ with 95% confidence interval.

RESULTS

We analyzed panoramic radiographs and charts of 1,001 orthodontic patients (45.8% males and 54.2% females). At least one dental anomaly was found in 34.8% ($n = 348$) of patients. The distribution of dental anomalies by sex showed that females were more affected than males (19.3% vs. 15.5%). The prevalence of investigated developmental dental anomalies of number, size, and position is presented in Figure 1. The location, number of teeth affected by DDA in the upper and lower jaw and comparison between males and females are presented in Tables 1 and 2.

Abnormalities of tooth number

Tooth agenesis was the most frequent abnormality of tooth number presented in 12.9% ($n = 129$) of all patients (5.5% of males and 7.4% of females). Supernumerary teeth, including mesiodens, were observed in 4.4% ($n = 44$) of subjects (2.4% of males and 2% of females). A total of 2.5% ($n = 25$) patients had mesiodens (2.8% males and 2.2% females). And other kinds of supernumerary teeth were reported in 2.4% of patients. Prevalence of abnormalities of tooth number in male and female orthodontic patients is presented in Figure 2. The most commonly missing tooth was upper left second premolar ($n = 46$ teeth), followed




Figure 1. Prevalence of developmental dental anomalies in Serbian orthodontic patients




Figure 2. Number of male and female patients with anomalies of tooth number




Figure 3. Number of male and female patients with anomalies of tooth position

by upper right and lower right second premolar (37 teeth in both right quadrants). In the anterior region of maxilla, lateral incisors showed the highest prevalence of agenesis ($n = 40$ teeth). We found 17 lateral incisors missing on the left side, and 23 on the right side of maxilla. In the anterior region of the lower jaw, agenesis of incisors was the most frequent finding (21 teeth). First molars were not affected by agenesis. More second molars were missing in the lower jaw compared to the upper jaw (22 vs. 15 teeth). Oligodontia was reported in one female patient (Table 1 and 2).

Abnormalities of tooth position

Tooth impaction was the most frequently found dental abnormality (16.5%). The number of male and female patients with anomalies of tooth position is presented in Figure 3. The high number of impacted canines in the upper arch is documented in current study (107 teeth). We found 49 impacted canines on the right side, and 58 on the left side. Bilaterally impacted canines were present in 24 patients. Only 11 mandibular canines were impacted (five on the right side and six on the left side). More impacted premolars were found in the lower jaw. The only statistically significant difference in the number of teeth affected by DDA between males and females was found in the number of impacted maxillary second molars ($p < 0.05$) (Tables 1 and 2).

Abnormalities of tooth size

The DDA affecting tooth size were present in 4.7% ($n = 47$) of all patients in the sample. Prevalence of microdontia and macrodontia in male and female orthodontic patients is presented in Figure 4. The location, prevalence, and sex distribution of teeth affected by an abnormality in size are presented in Tables 1 and 2.

DISCUSSION

The present study assessed sex distribution and prevalence of selected DDA in the sample of 1,001 orthodontic patients. Numerous studies presented epidemiological data




Figure 4. Number of male and female patients with anomalies of tooth size

and prevalence of DDA in either general population or pediatric and orthodontic patients. They vary in selection of methods, sample size, number of included anomalies, and results. The size of our sample was found to be either similar or larger in number of subjects, comparing to recently conducted investigations. Furthermore, the studies with the same purpose, conducted in different parts of the world, reported at least one dental anomaly in 5.4–45.7% of subjects [4, 14–17]. The prevalence of DDA in the present study was 34.8%, which may be because the sample consisted of patients referred to orthodontic treatment. Multifactorial etiology of dental anomalies, ethnical differences, and selection of DDA investigated in the study, inclusion and exclusion criteria contribute to the diversity of results.

Abnormalities of tooth number

Agenesis of one or more teeth could create malocclusions and esthetic and functional problems. Missing teeth were the most frequent abnormality of tooth number in the present sample of Serbian orthodontic patients (12.9%). The overall prevalence of missing teeth in recent studies was 0.027–21.6% [14, 16, 18–22]. Such considerable differences in results could be explained by variation in sample composition and size, ethnicity and methodology. In a systematic review of the literature, Rakhshan and Rakhshan [20] reported a significantly higher number of patients with tooth agenesis in the samples comprised of orthodontic patients, in comparison to the epidemiological samples and samples of dental patients. They argued higher prevalence of anomalies in patients seeking orthodontic treatment. The only two recent epidemiological studies in Serbia found a lower prevalence of missing teeth (6.28% and 5.34%, respectively) in comparison to our results [6, 7]. The high rate of hypodontia in Serbian orthodontic patients in a present study could be due to the nature of sample composition. Patients with the most challenging malocclusions, in need of potentially complicated and multidisciplinary treatment approach are almost automatically referred to the Clinic of Orthodontics. The present finding of more females than males with hypodontia (7.4% vs. 5.5%) supports the documented sex differences in the association between sex and hypodontia, microdontia,

Table 1. Location and prevalence of maxillary teeth affected by developmental dental anomalies. Comparison between males and females. χ^2 test and Fisher's exact test ($p < 0.05$)

Tooth	Sex	Hypodontia		Hyperdontia		p	Impaction		Transposition		p	Microdontia		Macrodontia		p
		n	(%)	n	(%)		n	(%)	n	(%)		n	(%)	n	(%)	
17	M	6	1.3	/		0.153	0	0	/		0.005*	/		/		/
	F	2	0.4				9	1.7								
16	M	/		/		/	/		/		/	/		/		/
	F															
15	M	21	4.6	3	0.7	0.332	6	1.6	/		0.796	/		/		/
	F	16	3	2	0.4		9	1.7								
14	M	1	0.2	/		0.382	/		1	0.2	1.000	/		/		/
	F	1	0.2						1	0.2						
13	M	1	0.2	/		1.000	25	5.5	1	0.2	0.658	/		/		/
	F	1	0.2				33	6.1	1	0.2						
12	M	8	1.7	3	0.7	0.549	1	0.2	/		1.000	6	1.3	1	0.2	0.873
	F	15	2.8	3	0.6		1	0.2				6	1.1	2	0.4	
11	M	0	0	/		1.000	10	2.2	/		0.477	/		2	0.4	0.120
	F	1	0.2				8	1.5						8	1.5	
21	M	0	0	/		0.256	6	1.3	/		0.526	/		4	0.9	0.761
	F	3	0.6				4	0.7						6	1.1	
22	M	7	1.5	3	0.7	0.468	2	0.4	/		0.209	7	1.5	1	0.2	0.310
	F	10	1.8	1	0.2		0	0				13	2.4	4	0.7	
23	M	3	0.7	/		1.000	22	4.8	2	0.4	0.894	/		/		/
	F	3	0.6				27	5	1	0.2						
24	M	1	0.2	/		1.000	/		2	0.4	1.000	/		/		/
	F	2	0.4						3	0.6						
25	M	19	4.1	1	0.2	0.571	6	1.3	/		1.000	/		/		/
	F	27	5	3	0.5		7	1.4								
26	M	/		/		/	/		/		1.000	/		/		/
	F															
27	M	5	1.1	/		0.257	0	0	/		0.256	/		/		/
	F	2	0.4				3	0.6								

M – males; F – females;

*statistically significant; χ^2 test

hyperdontia and macrodontia. Females are more affected by tooth agenesis and microdontia, while more supernumerary and large teeth are expected to be found in men (1:1.5 ratio) [1, 23]. Previous studies in different world regions offered conflicting results regarding sex distribution of patients with missing teeth [15, 16, 19]. However, our findings are in agreement with the results of sex distribution in the Serbian population (5.34–6.28%) [6, 7]. In addition, the location of teeth agenesis is in relationship to teeth position in morphogenic field, i.e. to the most distal tooth in the group affecting second premolars and lateral incisors, as well as third molars [1]. Thus, these teeth are frequently affected by agenesis (4.28–7.52%) which is in accordance with our results [4–7, 18]. A rare occurrence of oligodontia was reported in the Italian population (0.08%) which is in agreement with our result (0.09%) [24].

Contrary to the high prevalence of hypodontia, supernumerary teeth are less frequently found in healthy individuals (0.5–3.8%). The prevalence of supernumerary teeth in our sample was 4.4%, slightly higher than in recent studies [24, 25, 26]. The etiological pattern of sex distribution in association with supernumerary teeth is the opposite of hypodontia [1]. Males are more prone to the formation of supernumerary teeth than females, which is in agreement with our findings [24, 27]. Mesiodens was the most frequently detected

supernumerary tooth on panoramic X-rays in the present study. The lower prevalence of mesiodens was reported in Italian non-orthodontic subjects and French orthodontic patients (0.05% and 0.66%) [15, 24].

Impaction

The prevalence of patients with tooth impactions in the current study was high (16.5%) in comparison to the results of recent studies (2.6–7.1%) [15, 16, 28]. The highest rate of tooth impaction was found in maxillary canines, followed by maxillary central incisors, mandibular premolars and second molar. In the present study, unerupted maxillary incisors were more frequently found in female subjects compared to males, which is not in agreement with the previous report [29]. The prevalence of patients with impacted canines was 10.7% in the present study, in comparison to findings in the general population ranging from 0.6–8.4% [4, 16, 24]. The higher rate of impaction in the current study is probably due to the composition of the sample comprised of persons referred to orthodontic treatment. Females were more affected by impacted maxillary canines compared to males (9.6% vs. 6.9%). Patients, especially females, perceive missing tooth in the anterior region of maxilla as an aesthetic problem, which motivates them to seek out orthodontic treatment.

Table 2. Location and prevalence of mandibular teeth affected by developmental dental anomalies. Comparison between males and females. χ^2 test and Fisher's exact test ($p < 0.05$)

Tooth	Sex	Hypodontia		Hyperdontia		p	Impaction		Transposition		p	Microdontia		Macrodontia		p
		n	(%)	n	(%)		n	(%)	n	(%)		n	(%)	n	(%)	
37	M	8	1.7	/		0.153	4	0.9	/		0.538	/		/		/
	F	4	0.7				7	1.3								
36	M	/		/		/	/		/		/	/		/		/
	F															
35	M	21	4.6	3	0.7	0.332	6	1.3	/		0.796	/		/		/
	F	16	3	2	0.4		9	1.7								
34	M	/		/		/	/		1	0.2	0.457	/		/		/
	F								0	0						
33	M	/		/		/	4	0.9	1	0.2	0.517	/		/		/
	F						2	0.4	0	0						
32	M	2	0.4	/		0.596	/		/		1.000	6	1.3	1	0.2	0.873
	F	1	0.2									6	1.1	2	0.4	
31	M	6	1.3	0	0	0.297	/		/		/	0	0	/		1.000
	F	3	0.6	1	0.2							1	0.2			
41	M	2	0.4	1	0.2	0.539	/		/		/	0	0	/		1.000
	F	3	0.6	0	0							1	0.2			
42	M	3	0.7	0	0	0.330	/		/		/	/		/		/
	F	1	0.2	1	0.2											
43	M	/		/		/	2	0.4	/		0.535	/		/		/
	F						3	0.6								
44	M	/		/		/	/		/		/	/		/		/
	F															
45	M	18	3.9	1	0.2	0.268	6	1.3	/		1.000	/		/		/
	F	14	2.6	0	0		7	1.4								
46	M	/		/		/	/		/		1.000	/		/		/
	F															
47	M	6	1.3	/		0.536	5	1.1	/		1.000	/		/		/
	F	4	0.7				7	1.3								

M – males; F – females

Transposition

The maxillary canines and first premolars were found in complete transposition in four males and four females. Only a few recent studies reported a low prevalence of transposition (0.09%) which is in agreement with our findings (0.08%) [16, 30].

Microdontia and Macrodontia

Sogra et al. [16], found microdontia in 1.6% of Iranian orthodontic patients, while in a smaller sample, Baron et al. [15], reported 2.55%. Microdontia in Serbian patients was present in similar number of patients predominantly affecting maxillary lateral incisors. Low prevalence of macrodontia was reported in Iranian subjects (0.02%), which is in agreement with the results of our study (1.8%) [15].

Limitations

This study has a few limitations. Firstly, we assessed archived files of patients at the Clinic of orthodontics in 2016. The recruitment of patients could not be considered random since patients were already pre-selected from the general population and referred to orthodontic treatment.

Secondly, only charts and panoramic radiographs were used to evaluate DDA. That implied assessment of the sample for only selected DDA. In order to find the prevalence of all types of developmental dental anomalies, more extensive diagnostic methods should be included. Only DDA that can be observed with 100% accuracy on panoramic radiographs were reported making them more relevant than the findings of the rest. Third, DDA were reported in the sample of patients older than seven years of age. Dental abnormalities, such as impaction of canines and second molars, and agenesis of second premolars could not be observed in younger age groups. This could suggest a possible disparity in the diagnosis of DDA. Fourth, although, microdontia and macrodontia were evaluated by accepted reliable diagnostic method (visual examination and comparison), no additional confirmation was obtained from measurements on study models.

Despite the limitations, present findings could offer a foundation for much needed extensive epidemiological studies on DDA prevalence, sex distribution and association among different dental irregularities in the general population in Serbia and worldwide. Furthermore, this study provides information, which is of importance for dental practitioners.

CONCLUSIONS

The prevalence of developmental dental anomalies in Serbian orthodontic patients was 34.8%. At least one tooth anomaly was found in 15.5% of males, and 19.3% of females. The most frequently observed dental abnormality was tooth impaction, followed by tooth agenesis, hypodontia, and anomalies in tooth size and transposition. All investigated developmental dental anomalies, were more frequently present in females, except supernumerary teeth. The most commonly missing tooth was upper left premolar. The maxillary canines had the highest impaction rate. Mesiodens was the most frequently found supernumerary

tooth. The transposition of upper canine and first premolar was rare. The anomalies of tooth size predominantly affected incisors.

ACKNOWLEDGEMENTS

We thank Milos Selimir Stamenić (student at UBC School of Biomedical engineering program, Vancouver) for his contribution to the creation of this manuscript.

Conflict of interest: None declared.

REFERENCES

1. Brook AH. Multilevel complex interactions between genetic, epigenetic and environmental factors in the aetiology of anomalies of dental development. *Arch Oral Biol.* 2009; 54 Suppl 1:S3–17.
2. Thesleff I. The genetic basis of tooth development and dental defects. *A J Med Genet A.* 2006; 140(23):2530–5.
3. Berkiroglu N, Mete S, Ozbay G, Yalcinkaya S, Kargul B. Evaluation of panoramic radiographs taken from 1,056 Turkish children. *Niger J Clin Pract.* 2015; 18(1):8–12.
4. Dang HQ, Constantine S, Anderson PJ. The prevalence of dental anomalies in an Australian population. *Aust Dent J.* 2017; 62(2):161–4.
5. Gracco AL, Zanatta S, Forin Valvecchi F, Bignotti D, Perri A, Baciliero F. Prevalence of dental agenesis in a sample of Italian orthodontic patients: an epidemiological study. *Prog Orthod.* 2017; 18(1):33.
6. Jankulovski S, Filipović G. Hipodoncija stalnih zuba dece iz Knjaževca. *Journal of the Anthropological Society of Serbia.* 2009; 44:141–6.
7. Janosevic M, Filipovic G, Rusic-Tasic V. Frequency of hypodontia of permanent teeth in children. *Acta Stomatologica Naissi.* 2004; 20(45):219–26.
8. Markovic D, Petrovic B, Peric T. Case series: clinical findings and oral rehabilitation of patients with amelogenesis imperfecta. *European archives of paediatric dentistry : official journal of the European Academy of Paediatric Dentistry.* 2010; 11(4):201–8.
9. Martinovic B, Ivanovic M, Cvetkovic A, Todic J, Milojkovic Z, Pavlovic J, et al. Prevalence, characteristics and severity of hypomineralization of the first permanent molars and incisors in children from the northern part of Kosovo and Metohija. *Srp Arh Celok Lek.* 2017; 145(7–8):364–9.
10. Lam EWN. Dental anomalies. In: White WC, Pharoah MJ, editors. *Oral radiology principles and interpretations.* 7th ed. St. Louis: Mosby; 2014. p. 582–611.
11. Garib DG, Peck S, Gomes SC. Increased occurrence of dental anomalies associated with second-premolar agenesis. *Angle Orthod.* 2009; 79(3):436–41.
12. D'Souza R, Kapadia H, Vieira A. Teeth. In: Stevenson RE, Hall JG, editors. *Human malformations and related anomalies.* New York: Oxford University Press; 2006.
13. Becker A. *The Orthodontic Treatment of Impacted Teeth.* 2nd ed. London: Informa; 2007.
14. Chan GXL, Tan ELY, Chew MT, Wong HC, Foong KWC, Yow M. Secondary dentition characteristics in an ethnic Chinese orthodontic population: A retrospective cross-sectional study. *J Investig Clin Dent.* 2019; 10(3):e12421.
15. Baron C, Houchmand-Cuny M, Enkel B, Lopez-Cazaux S. Prevalence of dental anomalies in French orthodontic patients: A retrospective study. *Arch Pediatr.* 2018; 25(7):426–30.
16. Sogra Y, Mahdjoub GM, Elham K, Shore TM. Prevalence of dental anomalies in Iranian orthodontic patients. *Journal of Dentistry and Oral Hygiene.* 2012; 4(2):16–20.
17. Yassin SM. Prevalence and distribution of selected dental anomalies among Saudi children in Abha, Saudi Arabia. *J Clin Exp Dent.* 2016; 8(5):e485–e90.
18. Acev DP, Gjorgova J. Prevalence of Hypodontia in the Permanent Dentition of Macedonian Population. *Balkan J Dent Med.* 2014; 18(2):93–8.
19. Souza-Silva BN, Vieira WA, Bernardino ÍM, Batista MJ, Bittencourt MAV, Paranhos LR. Non-syndromic tooth agenesis patterns and their association with other dental anomalies: A retrospective study. *Arch Oral Biol.* 2018; 96:26–32.
20. Rakhshan V, Rakhshan H. Meta-analysis and systematic review of the number of non-syndromic congenitally missing permanent teeth per affected individual and its influencing factors. *Eur J Orthod.* 2016; 38(2):170–7.
21. Bandaru BK, Thankappan P, Kumar Nandan SR, Amudala R, Annem SK, Rajendra Santosh AB. The prevalence of developmental anomalies among school children in Southern district of Andhra Pradesh, India. *J Oral Maxillofac Pathol.* 2019; 23(1):160.
22. Hagiwara Y, Uehara T, Narita T, Tsutsumi H, Nakabayashi S, Araki M. Prevalence and distribution of anomalies of permanent dentition in 9,584 Japanese high school students. *Odontology.* 2016; 104(3):380–9.
23. Brook AH. A unifying aetiological explanation for anomalies of human tooth number and size. *Arch Oral Biol.* 1984; 29(5):373–8.
24. Lagana G, Venza N, Borzabadi-Farahani A, Fabi F, Danesi C, Cozza P. Dental anomalies: prevalence and associations between them in a large sample of non-orthodontic subjects, a cross-sectional study. *BMC oral health.* 2017; 17(1):62.
25. Fekonja A. Prevalence of dental developmental anomalies of permanent teeth in children and their influence on esthetics. *J Esthet Restor Dent.* 2017; 29(4):276–83.
26. Bilge NH, Yeşiltepe S, Törenek Ağırman K, Çağlayan F, Bilge OM. Investigation of prevalence of dental anomalies by using digital panoramic radiographs. *Folia Morphol (Warsz).* 2018; 77(2):323–8.
27. Bartolo A, Camilleri A, Camilleri S. Unerrupted incisors—characteristic features and associated anomalies. *Eur J Orthod.* 2010; 32(3):297–301.
28. Roslan AA, Rahman NA, Alam MK. Dental anomalies and their treatment modalities/planning in orthodontic patients. *J Orthod Sci.* 2018; 7:16.
29. Alhammadi MS, Asiri HA, Almashraqi AA. Incidence, severity and orthodontic treatment difficulty index of impacted canines in Saudi population. *J Clin Exp Dent.* 2018; 10(4):e327–e34.
30. Popoola B, Onyejaka N, Folayan MO. Prevalence of developmental dental hard-tissue anomalies and association with caries and oral hygiene status of children in Southwestern, Nigeria. *BMC oral health.* 2017; 17:8.

Заступљеност развојних анормалија зуба код ортодонтских пацијената у Србији

Евгенија Марковић¹, Ана Вуковић², Тамара Перић², Јована Кузмановић-Пфићер³, Бојан Петровић⁴

¹Универзитет у Београду, Стоматолошки факултет, Клиника за ортопедију вилица, Београд, Србија;

²Универзитет у Београду, Стоматолошки факултет, Клиника за дечју и превентивну стоматологију, Београд, Србија;

³Универзитет у Београду, Стоматолошки факултет, Катедра за медицинску статистику и информатику, Београд, Србија;

⁴Универзитет у Новом Саду, Клиника за стоматологију Војводине, Одељење дечје и превентивне стоматологије, Нови Сад, Србија

САЖЕТАК

Увод/Циљ Циљ овог истраживања је био да се испита заступљеност развојних анормалија зуба код ортодонтских пацијената у Србији.

Метод Узорак је чинио 1001 ортопантомографски снимак ортодонтских пацијената старијих од седам година са Клинике за ортопедију вилица Стоматолошког факултета у Београду. Бележено је присуство развојних анормалија за чију дијагностику је потребан само ортопантомографски снимак. За испитивање заступљености развојних анормалија зуба коришћена је дескриптивна статистичка анализа. χ^2 тест је коришћен ради поређења броја зуба са анормалијом између полова (степен значајности 95%).

Резултати Развојне анормалије зуба су биле заступљене код 34,8% ортодонтских пацијената (15,5% мушкараца и 19,3% жена). Импакције зуба су биле присутне код 16,5% пацијената, хиподонција код 12,9%, прекобројни зуби код 4,4%, микродонција код 2,9%, макродонција код 1,8% и транс-

позиција код 0,8% пацијената. Очњаци у горњој вилици су били најчешће импактирани зуби. Горњи други кутњаци су били склонији импакцији код жена ($p < 0,05$). Документовано је више импактираних секутића у горњој вилици, а преткутњака и других кутњака у доњој вилици. Најчешће су недостајали горњи леви преткутњаци. Од свих прекобројних зуба најчешће је био уочаван мезиоденс.

Закључак Приказали смо постојање високе учесталости развојних анормалија зуба код ортодонтских пацијената у Србији са већом израженошћу код особа женског пола. Најчешће анормалије биле су импакција, хиподонција, хипердонција, микродонција, макродонција и транспозиција. Све анормалије су биле учесталије код жена, осим у случају прекобројних зуба. Резултати садашње студије могу бити полазна тачка за епидемиолошке студије о учесталости развојних анормалија зуба.

Кључне речи: развојне анормалије зуба; ортодонција; хиподонција